

COX IV (PTR1399) Mouse mAb

CatalogNo: YM4289 **Recombinant** 

Key Features

Host Species

- Mouse

Reactivity

- Human, Mouse, Rat

Applications

- WB, IF, ELISA, IHC

MW

- 20kD (Calculated)
- 17kD (Observed)

Isotype

- IgG2b, kappa

Storage

Storage* -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

Recommended Dilution Ratios

IHC 1:200-1000**WB 1:500-2000****IF 1:100-500****ELISA 1:1000-5000**

Basic Information

Clonality Monoclonal**Clone Number** PTR1399

Immunogen Information

Immunogen AA range: 100-200**Specificity** This antibody detects endogenous levels of COX IV protein.

Target Information

Gene name COX4I1

Protein Name Cytochrome c oxidase subunit 4 isoform 1, mitochondrial


Organism	Gene ID	UniProt ID
Human	1327 ;	P13073 ;
Mouse	12857 ;	P19783 ;
Rat	29445 ;	P10888 ;

Cellular Localization Mitochondrion inner membrane


Tissue specificity Ubiquitous.

Function Function:This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport.,similarity:Belongs to the cytochrome c oxidase IV family.,tissue specificity:Ubiquitous.,


Validation Data




Immunogenic Radiation Therapy for Enhanced Antitumor Immunity via a Core-Shell Nanosensitizer-Mediated Immunosuppressive Tumor Microenvironment Modulation. ACS Nano Jin-Xiang Chen IF Mouse 4T1 cell-xenograft




Immunofluorescence analysis of HeLa cell. 1, Rabbit Histone H3 Antibody YM8335 (red) was diluted at 1:200 (4°C overnight).COX IV (PTR1399) Mouse mAb (green) was diluted at 1:200 (4°C overnight). 2, Goat Anti Rabbit Alexa Fluor 594 Catalog:RS3611 was diluted at 1:1000 (room temperature, 50min). Goat Anti Mouse Alexa Fluor 488 Catalog:RS3208 was diluted at 1:1000 (room temperature, 50min).




Immunofluorescence analysis of Mouse-kidney tissue. 1, COX IV Monoclonal Antibody (6C8) (red) was diluted at 1:200 (4°C overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B




Immunofluorescence analysis of Rat-kidney tissue. 1, COX IV Monoclonal Antibody (6C8) (red) was diluted at 1:200 (4°C overnight). 2, Cy3 labeled Secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-COX IV (PTR1399) antibody. The HRP-conjugated Goat anti-Mouse IgG (H + L) antibody was used to detect the antibody. Lane 1: MCF7 Lane 2: HEK293 Lane 3: 4T1 Lane 4: Rat heart Predicted band size: 20kDa Observed band size: 17kDa



Human kidney was stained with anti-COX IV (PTR1399) mouse antibody



Human liver was stained with anti-COX IV (PTR1399) Mouse antibody

Contact information

Orders: order@immunoway.com
 Support: tech@immunoway.com
 Telephone: 877-594-3616 (Toll Free), 408-747-0185
 Website: <http://www.immunoway.com>
 Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:
COX IV (PTR1399)
Mouse mAb

