

Goat Anti Rabbit IgG(H+L) (HRP)

CatalogNo: RS0002 Orthogonal Validated (

Key Features

Host Species

Goat

Reactivity

Applications Rabbit WB,IHC,ELISA

Conjugate

HRP

Recommended Dilution Ratios

WB 1:5000-20000 IHC 1:20-100

ELISA 1:5000-20000

Storage

Storage* -15°C to -25°C/1 year(Do not lower than -25°C)

Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.


I Basic Information

Immunogen Information


| Target Information

Protein Name


| Validation Data




TBBPA causes apoptosis in grass carp hepatocytes involving destroyed ERmitochondrial function. CHEMOSPHERE Shiwen Xu WB Fish 1:8000 L8824 cell



A novel CD8+ T cell-related gene signature for predicting the prognosis and immunotherapy efficacy in bladder cancer. INFLAMMATION RESEARCH Xu Ning WB Human 1:1000 5637 cell,T24 cell,UMUC-3 cell,SV-HUC-1 cell



Kang, Min, et al. "Autophagy was activated against the damages of placentas caused by nano-copper oral exposure." Ecotoxicology and Environmental Safety 220 (2021): 112364.



Western Blot analysis of flag-tag protein using primary antibody at various dilution. Secondary antibody(catalog#:RS0002) was diluted at 1:10000

| Contact information

Orders: order@immunoway.com
Support: tech@immunoway.com

Telephone: 877-594-3616 (Toll Free), 408-747-0185

Website: http://www.immunoway.com

Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:

Goat Anti Rabbit

IgG(H+L) (HRP)

For Research Use Only. Not for Use in Diagnostic Procedures.	Antibody ELISA Kits Protein Reagents
	Immunoway - 3 / 3