

## DDX58 (PT0686R) PT® Rabbit mAb

CatalogNo: YM8495 **Recombinant** 

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse

#### Applications

- WB, IHC, IF, IP, ELISA

#### MW

- 107kD (Calculated)
- 107kD (Observed)

#### Isotype

- IgG, Kappa

### Recommended Dilution Ratios

**IHC 1:200-1:1000****WB 1:500-1:2000****IF 1:200-1:1000****ELISA 1:5000-1:20000****IP 1:50-1:200**

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)**Formulation** PBS, 50% glycerol, 0.05% Proclin 300, 0.05% BSA

### Basic Information

**Clonality** Monoclonal**Clone Number** PT0686R

### Immunogen Information

**Specificity** Endogenous

---

## Target Information

**Gene name** DDX58

**Protein Name** Probable ATP-dependent RNA helicase DDX58 (DEAD box protein 58) (RIG-I-like receptor 1) (RLR-1) (Retinoic acid-inducible gene 1 protein) (RIG-1) (Retinoic acid-inducible gene 1 protein) (RIG-I)

Organism	Gene ID	UniProt ID
Human	<a href="#">23586;</a>	<a href="#">O95786;</a>
Mouse		<a href="#">Q6Q899;</a>


**Cellular Localization** Cytoplasm

**Tissue specificity** Present in vascular smooth cells (at protein level).


**Function** Domain:The 2 CARD domains are responsible for interaction with and signaling through MAVS.,Domain:The helicase domain is responsible for dsRNA recognition.,Domain:The repressor domain controls homomultimerization and interaction with MAVS.,Function:Involved in innate immune defense against viruses. Upon interaction with intracellular dsRNA produced during viral replication, triggers a transduction cascade involving MAVS/IPS1, which results in the activation of NF-kappa-B, IRF3 and IRF7 and the induction of the expression of antiviral cytokines such as IFN-beta and RANTES (CCL5). Essential for the production of interferons in response to RNA viruses including paramyxoviruses, influenza viruses, Japanese encephalitis virus and HCV.,induction:By bacterial lipopolysaccharide (LPS) in endothelial cells. By IFN-alpha, -beta and -gamma.,PTM:Isgylated. Conjugated to ubiquitin-like protein ISG15 upon IFN-beta stimulation.,similarity:Belongs to the helicase family.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 2 CARD domains.,subunit:Monomer; maintained as a monomer in an autoinhibited state. Upon viral dsRNA binding and conformation shift, homomultimerizes and interacts with MAVS. Interacts with DHX58/LGP2, IKBKE, TBK1 and TMEM173/STING.,tissue specificity:Present in vascular smooth cells (at protein level).,

---

## Validation Data



Human stomach was stained with anti-DDX58 (PT0686R) rabbit antibody



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-DDX58 (PT0686R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: Jurkat Lane 2: A549 Predicted band size: 107kDa Observed band size: 107kDa

## Contact information

Orders: [order@immunoway.com](mailto:order@immunoway.com)  
Support: [tech@immunoway.com](mailto:tech@immunoway.com)  
Telephone: 877-594-3616 (Toll Free), 408-747-0185  
Website: <http://www.immunoway.com>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**DDX58 (PT0686R)**  
**PT® Rabbit mAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)