

## MDA5 Rabbit pAb

CatalogNo: YT2687

### Key Features

#### Host Species

- Rabbit

#### Reactivity

- Human, Mouse

#### Applications

- WB, IHC, IF, ELISA

#### MW

- 120kD (Observed)

#### Isotype

- IgG

### Recommended Dilution Ratios

**WB 1:500-1:2000**

**IHC 1:100-1:300**

**ELISA 1:40000**

**IF 1:50-200**

### Storage

**Storage\*** -15°C to -25°C/1 year (Do not lower than -25°C)

**Formulation** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

### Basic Information

**Clonality** Polyclonal

### Immunogen Information

**Immunogen** The antiserum was produced against synthesized peptide derived from human IFIH1. AA range: 976-1025

**Specificity** MDA5 Polyclonal Antibody detects endogenous levels of MDA5 protein.

### Target Information

**Gene name** IFIH1

**Protein Name** Interferon-induced helicase C domain-containing protein 1


| Organism | Gene ID                | UniProt ID              |
|----------|------------------------|-------------------------|
| Human    | <a href="#">64135;</a> | <a href="#">Q9BYX4;</a> |
| Mouse    | <a href="#">71586;</a> | <a href="#">Q8R5F7;</a> |

**Cellular Localization** Cytoplasm . Nucleus . Mitochondrion . Upon viral RNA stimulation and ISGylation, translocates from cytosol to mitochondrion. May be found in the nucleus, during apoptosis.


**Tissue specificity** Widely expressed, at a low level. Expression is detected at slightly highest levels in placenta, pancreas and spleen and at barely levels in detectable brain, testis and lung.

**Function** Disease:Genetic variation in IFIH1 is associated with insulin-dependent diabetes mellitus 19 (IDDM19) [MIM:610155].,Function:RNA helicase that, through its ATP-dependent unwinding of RNA, may function to promote message degradation by specific RNases. Seems to have growth suppressive properties. Involved in innate immune defense against viruses. Upon interaction with intracellular dsRNA produced during viral replication, triggers a transduction cascade involving MAVS/IPS1, which results in the activation of NF-kappa-B, IRF3 and IRF7 and the induction of the expression of antiviral cytokines such as IFN-beta and RANTES (CCL5). ATPase activity is specifically induced by dsRNA. Essential for the production of interferons in response to picornaviruses.,induction:By IFN-beta and TNF-alpha.,miscellaneous:In HIV-1 infected HeLa-CD4 cells, overexpression of IFIH1 results in a great increase in the level of secreted viral p24 protein.,PTM:During apoptosis, processed into 3 cleavage products. The helicase-containing fragment, once liberated from the CARD domains, translocate from the cytoplasm to the nucleus. The processed protein significantly sensitizes cells to DNA degradation.,sequence Caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the helicase family.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 2 CARD domains.,subcellular location:May be found in the nucleus, during apoptosis.,subunit:Interacts with MAVS. Interacts with V protein of Simian virus 5, Human parainfluenza virus 2, Mumps virus, Sendai virus and Hendra virus. Binding to paramyxoviruses V proteins prevents IFN-beta induction, and the further establishment of an antiviral state.,tissue specificity:Widely expressed, at a low level. Expression is detected at slightly highest levels in placenta, pancreas and spleen and at barely levels in detectable brain, testis and lung.,

## Validation Data



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from Jurkat cells, using IFIH1 Antibody. The lane on the right is blocked with the synthesized peptide.

## Contact information

Orders: [order@immunoway.com](mailto:order@immunoway.com)  
Support: [tech@immunoway.com](mailto:tech@immunoway.com)  
Telephone: 877-594-3616 (Toll Free), 408-747-0185  
Website: <http://www.immunoway.com>  
Address: 2200 Ringwood Ave San Jose, CA 95131 USA



Please scan the QR code to access additional product information:  
**MDA5 Rabbit pAb**

For Research Use Only. Not for Use in Diagnostic Procedures.

[Antibody](#) | [ELISA Kits](#) | [Protein](#) | [Reagents](#)